

On the Twisted q -Euler numbers and polynomials associated with basic q - l -functions

TAEKYUN KIM, *EECS, Kyungpook National University, Daegu 702-701, S. Korea*
e-mail: tkim64@hanmail.net or tkim@knu.ac.kr

SEOG-HOON RIM, *Department of Mathematics Education, Kyungpook National University, Daegu 702-701, S. Korea*
e-mail: shrim@knu.ac.kr

ABSTRACT. One purpose of this paper is to construct twisted q -Euler numbers by using p -adic invariant integral on $\mathbb{Z}_p$ in the sense of fermionic. Finally, we consider twisted Euler q -zeta function and q - l -series which interpolate twisted q -Euler numbers and polynomials.

1. Introduction

Let p be a fixed odd positive integer. Throughout this paper $\mathbb{Z}_p$, $\mathbb{Q}_p$, $\mathbb{C}$ and $\mathbb{C}_p$ are respectively denoted as the ring of p -adic rational integers, the field of p -adic rational numbers, the complex numbers field and the completion of algebraic closure of $\mathbb{Q}_p$. The p -adic absolute value in $\mathbb{C}_p$ is normalized so that $|p|_p = 1/p$. When one talks of q -extension, q is considered in many ways such as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes that $|q| < 1$. If $q \in \mathbb{C}_p$, we normally assume that $|q - 1|_p < p^{-1/p-1}$ so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$, cf. [1, 2, 15, 20].

We use the notations as

$$\begin{aligned} [x]_q &= \frac{1 - q^x}{1 - q} = 1 + q + q^2 + \cdots + q^{x-1}, \\ [x]_{-q} &= \frac{1 - (-q)^x}{1 + q} \\ &= 1 - q + q^2 - q^3 + \cdots + (-1)^x q^{x-1}, \quad \text{see [12].} \end{aligned}$$

Let $UD(\mathbb{Z}_p)$ be the set of uniformly differentiable function on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the p -adic invariant q -integral on $\mathbb{Z}_p$ was defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{\mathbf{v} \rightarrow \infty} \frac{1}{[p^{\mathbf{v}}]_q} \sum_{x=0}^{p^{\mathbf{v}}-1} f(x) q^x, \quad (\text{see [3, 8, 9]}).$$

Note that

$$I_1(f) = \lim_{q \rightarrow 1} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_1(x) = \lim_{\mathbf{v} \rightarrow \infty} \frac{1}{p^{\mathbf{v}}} \sum_{x=0}^{p^{\mathbf{v}}-1} f(x).$$

For a fixed positive integer d with $(p, d) = 1$, set

$$\begin{aligned} X = X_d &= \varprojlim \mathbb{Z}/dp^{\mathbf{v}}\mathbb{Z} \\ X_1 &= \mathbb{Z}_p \\ X^* &= \bigcup_{0 < a < dp, (a, p) = 1} \\ a + dp^{\mathbf{v}}\mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^{\mathbf{v}}}\}, \end{aligned}$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^{\mathbf{v}}$, cf. [15].

Recently Kim consider q -Euler numbers as follows:

$$F_q(x, t) = [2]_q \sum_{n=0}^{\infty} (-1)^n q^n e^{\lceil n+x \rceil_q t} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}, \quad \text{see [5, 12].}$$

For $h \in \mathbb{Z}$, $k \in \mathbb{N}$, q -Euler polynomials of higher order were defined by

$$E_{n,q}^{(h,k)}(x) = \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{k\text{-times}} [x + x_1 + \cdots + x_k]_q^n q^{\sum_{i=1}^k x_i(h-i)} d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k), \quad \text{see [8, 15].} \quad (1)$$

The q -Euler polynomial at $x = 0$ are called q -Euler numbers, cf. [1, 2, 14]. Hence q -Euler numbers of higher order can be defined by $E_{n,q}^{(h,k)} = E_{n,q}^{(h,k)}(0)$.

Let χ be a Dirichlet Character with conductor $f(= \text{odd}) \in \mathbb{N}$. Then the l -function attached to χ is defined as

$$l(s, \chi) = 2 \sum_{n=1}^{\infty} \frac{\chi(n)(-1)^n}{n^s}, \quad \text{for } s \in \mathbb{C}, \quad \text{see [4, 13].}$$

When $\chi = 1$, this is the Euler zeta function, which is defined by

$$\zeta_E(s) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}, \quad \text{see [3, 4, 13].}$$

In [8, 12], q -analogue of Euler ζ -function is defined by

$$\zeta_{E,q}^{(h)}(s, x) = [2]_q \sum_{n=0}^{\infty} \frac{(-1)^n q^{nh}}{[n+x]_q^s}, \quad s, h \in \mathbb{C}. \quad (2)$$

Note that $\zeta_{E,q}^{(h)}(s, x)$ is analytic continuation an $\mathbb{C}$ with

$$\zeta_{E,q}^{(h)}(-n, x) = E_{n,q}^{(h,1)}(x), \quad \text{for } n \in \mathbb{N}, h \in \mathbb{Z}, \quad \text{see [8, 5].} \quad (3)$$

It was known that

$$E_{n,q}^{(h,1)}(x) = [2]_q \sum_{k=0}^{\infty} (-1)^k q^{kh} [k+x]_q^n, \quad \text{see [8]}. \quad (4)$$

It follows from (2) that

$$\lim_{q \rightarrow 1} \zeta_{E,q}^{(h)}(s, x) = \zeta_E(s, x) = 2 \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+x)^s}, \quad \text{see [13]}.$$

With meaning of q -analogue of Dirichlet's type l -function, we consider the following q - l -function : for $s \in \mathbb{C}$

$$l_q(s, \chi) = [2]_q \sum_{k=1}^{\infty} \frac{\chi(k) (-1)^k q^k}{[k]_q^s}, \quad \text{see [3, 5, 12]}. \quad (5)$$

It is easy to see that $l_q(s, \chi)$ is analytic continuation in whole complex plane.

The generalized q -Euler numbers attached to χ were also defined by

$$[2]_q \sum_{n=0}^{\infty} e^{[n]_q t} \chi(n) (-1)^n q^n = \sum_{n=0}^{\infty} E_{n,\chi,q} \frac{t^n}{n!}, \quad \text{see [5, 6, 12]}. \quad (6)$$

Note that

$$\int_{\mathbb{Z}_p} \chi(x) [x]_q^n d\mu_{-q}(x) = E_{n,\chi,q} \quad \text{and} \quad l_q(-n, x) = E_{n,\chi,q}, \quad \text{see [3, 5]}. \quad (7)$$

For $m \geq 0$, extended generalized q -Euler numbers attached to χ are defined as

$$\begin{aligned} E_{n,\chi,q}^{(h,k)} &= \underbrace{\int_X \cdots \int_X}_{k\text{-times}} \left[\sum_{l=1}^k x_l \right]_q^n q^{\sum_{j=1}^k (h-j)x_j} \left(\prod_{j=1}^k \chi(x_j) \right) d\mu_{-q}(x_1) \cdots d\mu_{-q}(x_k) \\ &= \frac{[2]_q^k}{[2]_{q^f}^k} [f]_q^n \sum_{a_1, \dots, a_k=0}^{f-1} q^{\sum_{l=1}^k (h-l+1)a_l} (-1)^{a_1+\dots+a_k} \\ &\quad E_{n,q^f}^{(h,k)} \left(\frac{a_1 + a_2 + \dots + a_k}{f} \right) \prod_{i=1}^k \chi(a_i), \quad (\text{see [8, 10]}). \end{aligned} \quad (8)$$

From (8), we derive

$$\begin{aligned} E_{n,\chi,q}^{(h,1)} &= \frac{[2]_q}{[2]_{q^f}} [f]_q^n \sum_{a=0}^{f-1} q^{ha} (-1)^a E_{n,q^f}^{(h,1)} \left(\frac{a}{f} \right) \chi(a) \\ &= [2]_q \sum_{k=1}^{\infty} \chi(k) (-1)^k q^{hk} [k]_q^n, \quad \text{see [5: p.5]}. \end{aligned} \quad (9)$$

From (9), we can consider the following q - l -function:

$$l_q^{(h)}(s, \chi) = [2]_q \sum_{k=1}^{\infty} \frac{\chi(k)(-1)^k q^{hk}}{[k]_q^s}, \quad \text{see [5, 8, 10],}$$

where $h, s \in \mathbb{C}$. Note that $l_q^{(h)}(-n, \chi) = E_{n, \chi, q}$, for $n \in \mathbb{N}$, $h \in \mathbb{Z}$.

In the present paper, we give twisted q -Euler numbers by using p -adic invariant q -integral on $\mathbb{Z}_p$ in the sense of fermionic. Moreover, we construct the q -analogues of Euler zeta function and q - l -function, which interpolate twisted q -Euler number and polynomials at negative integer.

2. Twisted q -extension of Euler numbers

In this section, we assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$. From the definition of p -adic invariant q -integral on $\mathbb{Z}_p$ in the sense of fermionic, we derive

$$\lim_{q \rightarrow -1} I_q(f_1) = I_{-1}(f_1) = -I_{-1}(f) + 2f(0), \quad \text{where } f_1(x) = f(x+1), \quad \text{see [7].} \quad (10)$$

Let $T_p = \cup_{n \geq 1} C_{p^n} = \lim_{n \rightarrow \infty} C_{p^n} = C_{p^\infty}$ be locally constant space, where $C_{p^n} = \{w \mid w^{p^n} = 1\}$ is the cyclic group of order p^n . For $w \in T_p$, we denote by $\phi_w : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ the locally constant function $x \mapsto w^x$.

If we take $f(x) = \phi_w(x)e^{tx}$, then we have

$$\int_{\mathbb{Z}_p} e^{tx} \phi_w(x) d\mu_{-1}(x) = \frac{2}{we^t + 1}, \quad \text{see [4, 11, 13, 16, 17, 18, 19].} \quad (11)$$

From (10), we derive

$$I_{-1}(f_n) = (-1)^n I_{-1}(f) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad \text{see [6, 7],} \quad (12)$$

where $f_n(x) = f(x+n)$. By (12), we easily see that

$$\int_X e^{tx} \phi_w(x) \chi(x) d\mu_{-1}(x) = \frac{2 \sum_{i=0}^{f-1} \chi(i) \phi_w(i) e^{it}}{w^f e^{ft} + 1}. \quad (13)$$

Now we define the analogue of Euler numbers as follows :

$$\frac{2 \sum_{i=0}^{f-1} \chi(i) \phi_w(i) e^{it}}{w^f e^{ft} + 1} = \sum_{n=0}^{\infty} E_{n, \chi, w} \frac{t^n}{n!}. \quad (14)$$

From (13) and (14), we note that

$$\int_X x^n \phi_w(x) \chi(x) d\mu_{-1}(x) = E_{n, \chi, w}. \quad (15)$$

In the viewpoint of (15), we consider twisted q -Euler numbers using p -adic invariant q -integral on $\mathbb{Z}_p$ in the sense of fermionic as follows :

$$E_{n,w,q}^{(h,1)}(x) = \int_{\mathbb{Z}_p} q^{(h-1)y} w^y [x+y]_q^n d\mu_{-q}(y). \quad (16)$$

Observe that $\lim_{q \rightarrow 1} E_{n,q,w}^{(h,1)}(x) = E_{n,w}(x)$. When $x = 0$, we write $E_{n,w}^{(h)}(0, q) = E_{n,w}^{(h)}$. Note that $E_{n,w,q}^{(h,1)}$ is the twisted form of $E_{n,q}^{(h,1)}$, see [8].

From (16), we derive

$$E_{n,w,q}^{(h,1)}(x) = \frac{[2]_q}{(1-q)^n} \sum_{j=0}^n \binom{n}{j} (-1)^j q^{xj} \frac{1}{1+q^{h+j}w} \quad (17)$$

Eq.(17) is equivalent to

$$E_{n,w,q}^{(h,1)}(x) = [2]_q \sum_{k=0}^{\infty} (-1)^k w^k q^{hk} [x+k]_q^n, \quad h \in \mathbb{Z}, n \in \mathbb{N}. \quad (18)$$

By (16), we see that

$$E_{n,w,q}^{(h,1)}(x) = \frac{[2]_q}{[2]} [d]_q^n \sum_{a=0}^{d-1} q^{ha} w^a (-1)^a E_{n,q^d}^{(h,1)}\left(\frac{x+a}{d}\right),$$

where $n, d(=\text{odd}) \in \mathbb{N}$.

Let χ be the Dirichlet character with conductor $f(=\text{odd}) \in \mathbb{N}$. Then we define the generalized twisted q -Euler numbers as follows : For $n \geq 0$,

$$E_{n,w,\chi,q}^{(h,1)} = \int_X \chi(x) q^{(h-1)x} w^x [x]_q^n d\mu_{-q}(x). \quad (19)$$

Note that $E_{n,1,\chi,q}^{(1,1)} = E_{n,\chi,q}$, see [3, 5, 12].

From (19), we can also derive

$$\begin{aligned} E_{n,w,\chi,q}^{(h,1)} &= [f]_q^n \frac{[2]_q}{[2]_{q^f}} \sum_{a=0}^{f-1} q^{ha} w^a \chi(a) (-1)^a E_{n,w^f,g^f}^{(h,1)}\left(\frac{a}{f}\right) \\ &= [2]_q \sum_{k=1}^{\infty} \chi(k) (-1)^k q^{hk} w^k [k]_q^n. \end{aligned} \quad (20)$$

It is east to check that

$$\lim_{q \rightarrow 1} E_{n,w,\chi,q}^{(h,1)} = E_{n,w,\chi}, \quad \text{see [4].}$$

3. q -Euler zeta function and q - l -functions

In this section, we assume that $q \in \mathbb{C}$ with $|q| < 1$. Here we construct twisted q -Euler zeta function and twisted q - l -function. Let $\mathbb{R}$ be the field of real numbers and let w be the p^r -th root of unity. For $q \in \mathbb{R}$ with $0 < q < 1$, $s \in \mathbb{C}$ and $h \in \mathbb{C}$, we define twisted Euler q -zeta function as follows :

$$\zeta_{E,q,w}^{(h,1)}(s) = [2]_q \sum_{k=1}^{\infty} \frac{(-1)^k w^k q^{hk}}{[x+k]_q^s}.$$

Note that $\zeta_{E,q,w}^{(h,1)}$ is an analytic continuation in $\mathbb{C}$. In [8], it is easy to see that

$$\zeta_{E,q,1}^{(h,1)}(s) = \zeta_{E,q}^{(h)}(s).$$

By (18), we easily see that

$$\zeta_{E,q,w}^{(h,1)}(-n) = E_{n,w,q}^{(h,1)}, \quad \text{for } n \in \mathbb{N}, h \in \mathbb{Z}.$$

We now also consider Hurwitz's type twisted q -Euler zeta function as follows : For $s \in \mathbb{C}$, define

$$\zeta_{E,q,w}^{(h,1)}(s, x) = [2]_q \sum_{k=0}^{\infty} \frac{(-1)^k q^{hk} w^k}{[x+k]_q^s}, \quad \text{for } s \in \mathbb{C}, h \in \mathbb{C}. \quad (21)$$

Note that $\zeta_{E,q,w}^{(h,1)}(s, x)$ has an analytic continuation on $\mathbb{C}$. By (18), we easily see that

$$\zeta_{E,q,w}^{(h,1)}(-n, x) = E_{n,w,q}^{(h,1)}(x), \quad \text{for } n \in \mathbb{N}, h \in \mathbb{Z}.$$

From [8], we note that

$$\zeta_{E,q,1}^{(h,1)}(s, x) = \zeta_{E,q}^{(h)}(s, x).$$

Let χ be Dirichlet's character with conductor $f(= \text{odd}) \in \mathbb{N}$. Then we define twisted q - l -function which interpolates twisted generalized q -Euler numbers attached to χ as follows : For $s \in \mathbb{C}$, $h \in \mathbb{C}$, define

$$l_{q,w}^{(h,1)}(s, \chi) = [2]_q \sum_{k=1}^{\infty} \frac{\chi(k) (-1)^k q^{hk} w^k}{[k]_q^s}. \quad (22)$$

For any positive integer n , we obtain

$$l_{q,w}^{(h,1)}(-n, \chi) = E_{n,w,\chi,q}^{(h,1)}, \quad \text{for } n \in \mathbb{N}, h \in \mathbb{Z}.$$

From (22), we can also derive

$$\begin{aligned} l_{q,w}^{(h,1)}(s, \chi) &= [2]_q \sum_{k=1}^{\infty} \frac{\chi(k) (-1)^k q^{hk} w^k}{[k]_q^s} \\ &= [f]_q^{-s} \frac{[2]_q}{[2]_{q^f}} \sum_{a=1}^f \chi(a) (-1)^a q^{ha} w^a \zeta_{E,q^f,w^f}(s, \frac{a}{f}). \end{aligned}$$

Question. Find a q -analogue of the p -adic twisted l -function which interpolates generalized twisted q -Euler numbers attached to $\chi E_{n,w,\chi,q}^{(h,1)}$ [4, 5, 10, 13].

References

- [1] L. Carlitz, q -Bernoulli and Eulerian numbers, Trans. Amer. Math. Soc., **76**(1954), 332-350.
- [2] M. Cenkci and M. Can, *Some results on q -analogue of the Lerch zeta function*, Advan. Stud. Contemp. Math., **12**(2006), 213-222.
- [3] T. Kim, *A note on q -Volkenborn integration*, proc. Jangjeon Math. Soc, **8**(2005), 13-17.
- [4] T. Kim, *On the analogs of Euler numbers and polynomials associated with p -adic q -integral on $\mathbb{Z}_p$ at $q = -1$* , J. Math. Anal. Appl. (2006), doi:10.1016/j.jmaa.2006.09.27
- [5] T. Kim, *On p -adic q - l -functions and sums of powers*, J. Math. Anal. Appl. (2006), doi:10.1016/j.jmaa.2006.07.071.
- [6] T. Kim, *A note on q -Euler numbers and polynomials*, math. NT/0608649.
- [7] T. Kim, *An invariant p -adic q -integral on $\mathbb{Z}_p$* , Appl. Math. Letters(accepted: see math. NT/0607792).
- [8] T. Kim, *q -Volkenborn integration and its applications*, math. NT/0510524.
- [9] T. Kim, *q -Volkenborn integration*, Russian J. Math. Phys., **9**(2002), 288-299.
- [10] T. Kim, L. C. Jang, S.-H. Rim and H.-K. Pak, *On the twisted q -zeta functions and q -Bernoulli polynomials*, Far East J. Appl. Math., **13**(2003), 13-21.
- [11] T. Kim, *On Euler-Barnes multiple zeta functions*, Russian J. Math. Phys., **10**(2003), 261-267.
- [12] T. Kim, *On the q -extension of Euler and Genocchi numbers*, J. Math. Anal. Appl., **326**(2007), 1458-1465.
- [13] T. Kim and S.-H. Rim, *Explicit p -adic expansion for alternating sums of powers*, J. Math. Anal. Appl. (In press).
- [14] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integral and Basic q -zeta function*, Trends in Mathematics (Information Center for Mathematical Sciences), **9**(2006), 7-12.
- [15] S.-H. Rim et al, *On the analogs of Bernoulli numbers and related identities and zeta and L -functions*, J. Korean Math Soc. (In press)
- [16] Y. Simsek, *Theorems on twisted L -function and twisted Bernoulli numbers*, Advan. Stud. Contemp. Math., **11**(2005), 205-218.
- [17] Y. Simsek, *Twisted (h,q) -Bernoulli numbers and polynomials related to twisted (h,q) -zeta function and L -function*, J. Math. Anal. Appl., **324**(2006), 790-804.
- [18] Y. Simsek, *On p -Adic Twisted q - L -Functions Related to Generalized Twisted Bernoulli Numbers*, Russian J. Math. Phys., **13**(3)(2006), 340-348.

- [19] Y. Simsek, *q-analogue of twisted l-series and q-twisted Euler numbers* , J. Number Theory, **110** (2005),267-278.
- [20] K. Shiratani, *On a p-adic interpolating function for the Euler numbers and its derivatives*, Mem. Fac. Sci. Kyushu Univ. Math., **39**(1985), 113-125.